

Using Virtual Reality to Help Reduce Anxiety: A Pilot Study

Anusha Devanga¹, Armili Paturi², Ganachari Vydyam Manogna³, Gauri Menon⁴ and Adithya Balasubramanyam⁵

¹⁻⁵CAVE Labs, PES University, Bangalore, Karnataka, India

Email: adithyab@pes.edu

Abstract— Anxiety is a common mental health condition that can be defined as a complex emotional state characterised by feelings of tension and worried thoughts. There are various treatment options available for anxiety, such as medication and therapy. However, recent research has explored the use of virtual reality in managing anxiety.

The Virtual Reality simulation acts as a channel for existing therapeutic methods and helps reduce the anxiousness they are experiencing. In our project, we have created two applications for the subjects to walk through. The applications incorporate principles of distraction therapy and guided meditation. Anxiety levels were assessed through continuous monitoring of the subject's blood pressure and heart rate during testing. A total of 37 subjects, aged between 16 and 40, participated in the testing of these applications. Through our results we observe that virtual reality could be a reliable channel for delivering therapeutic techniques to reduce anxiety.

Index Terms— Anxiety, Distraction therapy, Guided Meditation, Virtual Reality and Immersive Environment

I. INTRODUCTION

In stressful environments, it's common to feel nervous, but certain factors can escalate this feeling, causing stress, panic, and discomfort. The body releases hormones like adrenaline, triggering the 'fight or flight' response. However, some individuals may become overwhelmed, leading to anxiety. Although there are pharmacological methods [30] to treat anxiety, these medicines can have side effects. Virtual reality provides a safe environment, offering comfort and acting as a channel for therapeutic methods to alleviate anxiety. Our study aims to reduce anxiety in unavoidable stressful situations, like students taking exams, using virtual reality as a tool for existing therapeutic approaches.

A. Anxiety and its effects

Anxiety is characterised as an intense and persistent fear or worry. While it is completely normal to be nervous, anxiety is a more severe emotional state that can disrupt daily functioning [2] and lead to mental health issues. Symptoms of anxiety include excessive worrying, fear, restlessness, insomnia [1], sweating, trembling [26], increased heart rate, breathlessness [3], and elevated blood pressure [4].

Anxiety disorders refer to a group of mental disorders characterised by feelings of anxiety and fear, including generalised anxiety disorder (GAD) [37], panic disorder [38], phobias, social anxiety disorder [40], obsessive-compulsive disorder (OCD)[36] and post-traumatic stress disorder (PTSD)[39].

We intend to quantify the level of anxiety by measuring parameters that change when a person is anxious. Further

information regarding the parameters considered is provided in Section Parameters and rating system.

B. Virtual and Augmented Reality

The field of Virtual [6] and Augmented Reality (AR)[5] is gaining prominence due to its ability to offer personalised, immersive experiences that are easily accessible. It has proved to be useful in the automotive industry, the educational field as well as in the medical field [27]. The use of VR was found to be effective in reducing acute pain during and after different medical procedures. By providing an immersive environment that distracted patients from the source of their pain, VR was able to provide temporary relief [15].

Based on our research, we have developed and tested two applications incorporating distraction therapy and guided meditation. Anxiety levels were assessed through continuous monitoring of participants' physiological responses. The applications were tested on 37 subjects and demonstrated the efficacy of virtual reality as a reliable channel for delivering anxiety-reducing therapeutic techniques.

II. BACKGROUND

This section examines the existing applications that claim to help anxiety, therapy techniques, and the scales to measure anxiety. Additionally, the section provides a comprehensive review of prior research on the use of VR in addressing anxiety.

A. Existing Games to Reduce General Anxiety

Using games and other such applications to reduce anxiety is not a new concept. "SuperBetter" [10] is one such game. This application allows the user to 'fight' 'bad guys' which are certain habits that the user is trying to break. The application provides a daily set of challenges that the user needs to follow in real life which, on completion, provides a 'daily dose of power-ups'.

"Flower" [11] is another game meant to help reduce anxiety. It relies entirely on visual representation to act as a distraction for the user and does not have any text or dialog. The objective of the game is to distract the user long enough to calm them down.

Another application that was developed to help with users' mental health is "Personal Zen" [12]. The user is presented with a questionnaire and a personalised set of goals is given based on their answers which determine their stress levels.

B. Therapy Techniques

While there are various types of medication available to reduce anxiety, we focus on non-pharmacological approaches to do the same. One such method is distraction therapy, which aims to reduce anxiety by encouraging a subject to turn their attention to something other than the ongoing situation [13]. Another approach is meditation, which can promote relaxation and focus by directing the subject's attention to calming stimuli and eliminating stressful thoughts [24]. Guided meditation [23] is a form of meditation integrated into psychotherapy, that involves a guide leading the subject through the meditation session using various techniques such as soothing language, visualisations, or breathing exercises.

However, traditional therapy methods like distraction and guided meditation have limitations, such as requiring a calm and quiet environment that may not always be available. In our study, we developed VR applications to tackle the shortcomings of traditional therapy methods by creating a therapeutic and tranquil environment for subjects.

C. Existing Virtual Reality Applications for Anxiety

There are VR applications that exist in various industries. In this section, we will discuss a few examples. Pre-operative Anxiety: Preoperative Anxiety is an anxiety due to fear of medical procedures, mostly surgery or operations. In a research study, a group of patients were put through a Virtual Operating Room Tour (VORT) before administering anaesthesia [8][14].

Public Speaking Anxiety: The use of Virtual Reality Therapy (VRT) was employed to alleviate public speaking anxiety among university students, which is a type of social phobia [7].

Other non-pharmacological interventions such as cognitive-behavioural therapy, music therapy, aromatherapy, hypnosis, guided imagery relaxation therapy are becoming popular treatments for anxiety [9]. Augmented and virtual reality is currently being used to bring these concepts to life.

D. Existing scales of Anxiety

Assessing a subject's anxiety levels typically involves measures that are administered by self report[16]. Beck Anxiety Inventory (BAI)[16], State-Trait Anxiety Index (STAI)[16], and Hospital Anxiety and Depression Scale-Anxiety (HADS-A) [16] are the scales used to measure general anxiety. The STAI is among the most widely

researched and widely used measures of general anxiety and chronic medical conditions [16]. The BAI is used as a measure to differentiate between anxiety and depression [17], with increasing use in rheumatic conditions such as fibromyalgia [18] and arthritis [19]. The HADS-A scale is used in subjects to identify anxiety and depression symptoms. It includes an assessment of generalised anxiety including tension, worry, fear, panic, difficulties in relaxing, and restlessness [16].

III. METHODOLOGY

We classify anxiety levels into broad categories based on heart rate as shown in Table I. The values range from 60-220. These values are scaled down for classification, drawing inspiration from the Borg Rating of Perceived Exertion (RPE) scale [28]. First, we initialise the range from 0. So the new range will be 0-160. Dividing this by 10 will give us a new scale ranging from 0-16. Using this scale, subjects are classified into one of three categories. This predetermined scale is used only once - to determine which game needs to be administered.

TABLE I. CLASSIFICATION OF SUBJECTS

Class No.	Scale	Reading
0	0 - 5	Normal
1	5 - 10	Mild to Moderate Anxiety
2	10 - 15	Severe Anxiety

Based on this scale, two games were developed; each tailored to a specific class of anxiety. Subjects belonging to Class 1 were provided with a PC game. For subjects categorised under Class 2, a fully immersive VR environment was utilised. Details of each game are explained in this section.

The decision to opt for a PC game for Game 1, as opposed to a VR application like Game 2, was driven by the inherent complexity of the former, which demanded a higher level of user interaction facilitated by advanced input devices, such as a keyboard and a mouse. This augmented level of user interaction, in turn, implies that the application will hold the subject's attention and will be able to facilitate mild to moderate level anxiety.

A. Game 1: PC Game

Game 1 is a computer-based first-person movement game that uses the principles of distraction therapy. The game prompts subjects to solve puzzles and locate hidden treasures, engage them in the game world and redirect their attention away from their anxiety. At the start of the game, the subject is placed in a small coastal town and given the task of uncovering hidden "keys" that are scattered throughout the environment. These keys are three-dimensional objects - a cube, a prism, and an octahedron - that the subject must locate. When a key is found, a larger version of the same shape is created at the subject's original spawn point, with only the vertices visible. The subject must then return to the spawn location and complete the faces of this 3D model using triangles, which can be generated by selecting any three vertices in a clockwise direction. The selected vertices glow to indicate the same. If a triangle already exists between the selected vertices, it is erased. Once all the faces of the model are completed, the subject can move on to find the next key.

To navigate the town, players are provided with instructions at the start of the game as shown in Fig. 1. The controls are similar to those used in other first-person PC games, with walking controlled by the arrow keys or the keys 'W, A, S, D', and running achieved by pressing and holding the 'shift' button. Subjects can look around by clicking and dragging the mouse, simulating the motion of turning their heads.

B. Game 2: VR Game

Game 2 offers a fully immersive virtual environment designed for relaxation and it uses distraction therapy, as well as guided meditation. To access this virtual world, subjects are provided with a Head-mounted display (HMD) and controllers. Once in the virtual zen garden, they can use the controllers to teleport to different predetermined locations and explore the environment freely. A virtual pet dog and butterfly, along with coloured flares, provide additional features that follow the subject's desired path. The game includes a deep breathing exercise that the subject must follow as shown in Fig. 2. The instructions appear in the sky and include a voice-over that asks the subject to inhale, hold their breath, and exhale for a set duration, helping them to take deep breaths and relax. By immersing the subject in this environment and guiding them through meditation, the game aims to remove them from an anxiety-inducing environment and put them in a more relaxing setting. Game 2 only has two controls. The trigger button on the right-hand controller acts as a teleport button, enabling the subject to move to different predetermined locations. The touchpad on the left-hand controller allows for free movement in the environment,

enabling the subject to navigate by holding the touchpad in the direction they want to move. There is no particular objective or end goal in this game. The subject is free to explore the virtual world at their own pace, take in the scenery, and relax.




Fig 1. (a) Home Screen of the Game. (b) Key in the form of a Prism to trigger the vertices in the shape of a Prism. (c) When a vertex is selected it glows. The first vertex selected is blue and the second is green. (d) The subject is given instructions on player movement at the start of the game




Fig 2. (a) Virtual zen garden complete with cherry blossom trees and a tree-house. (b) Coloured flares and a butterfly follow the subject's right-hand movements. (c) One of the many comfort dogs that are available throughout the environment. (d) Guided meditation instructing the subject to Inhale. The bubble expands when the command is 'inhale'.

IV. TESTING

A. Technology and Hardware

a. Unity

The development of this project was done using Unity. Unity's cross-platform functionality aids in building applications that can be run on multiple platforms like Windows, Android, IOS, etc. Additionally, Unity offers development features specifically designed for VR programming, such as multi-platform support for various headsets, high-quality graphics with flexible graphic pipelines like Universal Render Pipeline and High Definition Render Pipeline, and advanced performance tools. These capabilities make Unity an ideal platform for VR development [20][21]

b. Virtual Reality System

HTC VIVE Cosmos Elite was chosen as the VR system. It comprises 2 VIVE Base Station 1.0, Cosmos Elite Headset and 2 VIVE controllers. Room-scale of approximately 11' 5" x 11' 5" is required for the tracking area. The Base Station 1.0 generates accurate external tracking. The VR setup used for this project is shown in Fig. 3.




Figure 3. The VR setup that was established for this project

SteamVR is a platform for virtual reality that includes a software framework designed to allow subjects to experience VR content and games on supported VR headsets. It also functions as a VR platform, enabling users to access and play VR games and content. The VIVE, which is compatible with SteamVR, offers freedom of movement and precision from all angles with SteamVR tracking and advanced controllers [22].

A motion faceplate is affixed to the front of a VR headset to enhance its motion tracking capabilities by detecting head movement and rotation. Additionally, the VIVE features on-ear, form-fitting headphones with integrated stereo audio to enhance the immersive experience. The VR System also includes several inputs such as a Multifunction trackpad, Dual stage trigger, Grip buttons, and an Integrated microphone. With a combined pixel resolution of 2880x1700, users can appreciate finer details within the VR application [25].

c. Parameters and Rating System

Prior research [31][32] has shown that anxiety disorders can have an effect on the Heart Rate Variance (HRV) of an individual. Individuals with anxiety had significantly lower resting-state HRV than the healthy population. Further, it was also found that HRV is inversely related to heart rate [33].

Research has shown that anxiety can increase heart rate, systolic blood pressure, and can cause an increased mean arterial pressure [34]. Considering this information as well as available equipment, heart rate, and blood pressure were deemed suitable indicators for determining the presence of anxiety in the subjects under examination.

The subjects were administered the State-Trait Anxiety Inventory (STAI) test. Form Y1 [29] was given to assess the subject's current state of anxiety. The scores in this test can range from 20 to 80. A cut point of 39 - 40 has been suggested to detect clinically significant symptoms [16]. For the purposes of this study, the sample consisted of subjects who scored 39 or above on the STAI test.

The subjects were varied across many categories: age group, sex, gaming experience and whether they had anxiety in general. The subject's heart rate in bpm and blood pressure was monitored and noted during the administration of the therapy technique supported by the proposed application. Subjects were asked to wear two smartwatches, one on each hand, and the readings were noted via blue-tooth on the watch apps.

The results were classified as positive or negative. If the value of the heart rate after the testing is lower than before, then it is a positive result. For blood pressure, non-escalation in value is a positive result. To further analyse the results, a rating system was also introduced. Heart rate and blood pressure are the two parameters for this experiment and the rating is done separately for both of them as shown in Table II and III.

TABLE II. THE RATING SYSTEM FOR HEART RATE AS A PARAMETER.

Heart Rate Value Range	Category	Rating
60-100	Normal	0
100-140	Low	1
140-180	Medium	2
180-220	High	3

TABLE III. THE RATING SYSTEM FOR BLOOD PRESSURE AS A PARAMETER.

Systolic (Upper\#)		Diastolic (Lower\#)	Blood Pressure Category	Rating
Less than 90	and	Less than 60	Low	0
Less than 120	and	Less than 80	Normal	1
120 - 129	and	Less than 80	Elevated	2
130 - 139	or	80 - 89	Higher Stage 1	3
140 or higher	or	90 or higher	Higher Stage 2	4

The aim of the application is to see a reduction in these parameters of the subject. Each subject is first classified into one of the categories of the tables (Table II and III). After the application is administered, they are classified again. If they remain in the same category, or they fall down a category, it is considered a 'positive' result. If they are in a higher category than before, it is considered a 'negative' result.

As the rating system is based on intervals, individual values of heart rate were not of significance within this range by itself. For example, let us assume the heart rate was 76 bpm before the application was administered and it fell to 86 bpm after. Even though the heart rate has increased, it is still in the same interval and is not considered a negative result. The applications, put together, were tested on a total of 37 subjects.

d) Testing Process

In this study, heart rate and blood pressure measurements were taken throughout each application. Measurements were recorded before the application was administered, three times during the testing, and once after the testing. The level of gaming experience that a subject has may impact the test results, and therefore, subjects were categorised into two groups: those with gaming experience (G) and those without gaming experience (NG). The criteria for this categorization was based on the subject's familiarity with games that involve First Person Movement (FPM) controls, as these are typically easier to understand for individuals with gaming experience. Subjects were queried about whether they had any prior experience with anxiety (HA) or not (NA). Table IV provides a comprehensive overview of the number of subjects who participated in the testing of the applications, along with their corresponding classifications into the predetermined categories.

V. RESULTS

A. Game 1

In regards to heart rate rating (based on intervals), all 14 participants exhibited a 100% positive result. As for blood pressure rating, there was a 93% positive outcome, with only one participant experiencing a slightly elevated blood pressure.

The graph in Fig. 4 displays the variation in the subjects' heart rates during the different phases of the application (as explained in section Game 2). The data reveals a noticeable decline in the average heart rate during the initial stages, followed by a gradual increase over time. Nevertheless, the average heart rate remains slightly lower after the game is administered as compared to the level before.

a) Gender

Gender did not play any part in influencing the result.

b) Gaming Experience

Considering the age of the test subjects, gaming experience could be considered prominent. But the results showed no significant difference. The subjects with no gaming experience have all shown positive results in both heart rate rating and blood pressure rating.

c) Presence of Anxiety

4 subjects reported to have experienced some form of anxiety and 3 of them showed positive reactions to the application.

TABLE IV. SUMMARY OF TEST SUBJECTS FOR ALL APPLICATIONS

Characteristic	Game 1	Game 2
Number of Subjects	14	23
Age (in years)	16-30	(18-40) G1 = 18 - 30 G2 = 30 - 40
Gender	9 Male, 5 Female	17 Male, 6 Female
Presence of Anxiety	4 HA, 10 NA	7 HA, 16 NA
Gaming Experience	9 G, 5 NG	8 G, 15 NG
Time Taken	No Time Limit	Controlled Simulation

d) Time Taken

Considering the age of the test subjects, gaming experience could be considered prominent. But the results showed no significant difference. The subjects with no gaming experience have all shown positive results in both heart rate rating and blood pressure rating.




Figure 4. Variation in heart rate for subjects who tested Game 1. The black line indicates the average heart rate of all the subjects.

B. Game 2

All 23 subjects showed a 100% positive result in the case of heart rate rating (based on intervals). The rating remained the same for 21 subjects while it was reduced for 2 subjects. Even individual heart rate values showed 95.65% of positive results with 22 out of 23 subjects experiencing lower heart rates after the simulation. For the subject with the exception, the heart rate value had only increased by 1 unit, leading to the same rating, nevertheless being a positive result.

The blood pressure rating also had a positive result. 78.3% of the subjects had either a consistent rating or a reduced rating.

The presented graph in Fig. 5 illustrates the variation in the subjects' heart rates while immersed in the virtual environment (see section Game 2). The data highlights a consistently slight decrease in the average heart rate as the subjects progress through the simulation.

a) Age

While the heart rating remains positive among both age groups, the blood pressure rating had a few exceptions. 78% of the subjects in Age Group 1 showed a decrease in heart rate while Age Group 2 had 67%.

b) Gender

Even if the rating for heart rate did not really show the influence of gender, it is interesting to see its variation in the results of blood pressure rating. 87.5% of the male subjects showed a decrease in heart rate while 67% of the female subjects showed a decrease in heart rate.

c) Gaming Experience

Even though 8 of them had prior experience, the nature of the result was not influenced, in fact, it was the opposite. The group with no gaming experience had a higher positive result than the one with gaming experience.

d) Presence of Anxiety

The heart rate value and rating had 100% positive results on subjects with and without anxiety.

e) Time Taken

It was found that the simulation time varied from person to person since each subject took different times to adjust to the environment but in totality the simulation lasted for approximately 8 minutes.




Figure 5. Variation in heart rate for subjects who tested Game 2. The black line indicates the average heart rate of all the subjects.

C. Analysis

Despite a slight increase in their heart rates and blood pressure, all subjects remained in the same category as mentioned in Table II and III, resulting in a 100% positive outcome for each of them. Taking decreased heart rate value into consideration, the results are 64.3%, and 95.65% positive for Game 1 and 2 respectively while for Blood pressure, the results are 93% and 78% positive.

Since subjects of both the applications saw a very slight variation in heart rate before and after the application was administered, merely reporting an increase or decrease in heart rate lacks sufficient detail without analysing the exact magnitude of change.

It is essential to examine the precise difference in heart rate. Table V displays the number of participants with an increased or decreased heart rate and the average bpm change.

The tabulated data indicate that Game 1 was successful in regulating the variability in heart rate, as the positive and negative impacts of the applications seemed to offset each other. However, Game 2, which was a VR game, stood out for its exceptional performance in reducing heart rate, suggesting that it effectively managed anxiety levels.

TABLE V. THE QUANTIFIED VALUES OF THE POSITIVE AND NEGATIVE IMPACT OF THE TWO APPLICATIONS ON HEART RATE.

	Game 1	Game 2
No. of subjects with a positive result for heart rate comparison	9	22
The average decrease in heart rate (in bpm)	8.11	10.36
No. of subjects with a negative result for heart rate comparison	5	1
The average increase in heart rate (in bpm)	7.4	1

VI. DISCUSSIONS

The applications are not intended to provide any form of behavioural therapy. Rather, their primary aim is to address anxiety by providing immediate relief to subjects. The design of the applications is geared towards delivering a short-term solution to anxiety, and therefore, the study's focus is on assessing its short-term effects on the subjects. Although our research did not consider any experiments designed to evaluate how the therapy would impact the subjects' well-being in the long term, we can make an informed guess as to whether it should either have a positive impact or surely shouldn't worsen the situation.

VR, although incredibly useful, has a few drawbacks. Prolonged exposure to immersive environments and the use of headsets have many side effects such as nausea, dizziness, eye strain, headache, and motion sickness. The applications also require the subject to be of sound mind and in the ability to move their hands and head.

VII. CONCLUSIONS

In this study, two distinct games were developed to act as viable channels for distraction therapy and guided meditation. The results have shown the effectiveness of VR in reducing anxiety. However, it is important to note that this is only a pilot study and doesn't account for specific phobias or triggers that an individual may have. For example, Game 2 features virtual dogs that could cause discomfort for some participants. Future research could focus on increasing the customization of the virtual environments to provide the best possible calming setting for each individual.

It's important to note that while the games provide short-term relief, they don't offer a long-term solution for underlying anxiety issues. To enhance these applications, incorporating Artificial Intelligence and integrating other forms of therapy can be considered. We remain fascinated by the concept of virtual reality and recognize it as a stepping stone towards making a difference in the future.

ACKNOWLEDGMENT

The work was done in partial fulfilment of the requirements for the bachelor's degree in computer science and engineering. We would like to thank PES University for providing us the required resources, place to work and funds for the current work.

REFERENCES

- [1] U.S. Department of Health and Human Services. (n.d.). Anxiety disorders. National Institute of Mental Health. Retrieved March 30, 2023, from <https://www.nimh.nih.gov/health/topics/anxiety-disorders>
- [2] World Health Organization. Depression and other common mental disorders: global health estimates. No. WHO/MSD/MER/2017.2. World Health Organization, 2017.
- [3] Currow, David C., Sungwon Chang, Helen K. Reddel, Slavica Kochovska, Diana Ferreira, Irina Kinchin, Miriam Johnson, and Magnus Ekström. "Breathlessness, anxiety, depression, and function—the BAD-F study: a cross-sectional and population prevalence study in adults." *Journal of pain and symptom management* 59, no. 2 (2020): 197-205.
- [4] Gorman, Jack M., and Richard P. Sloan. "Heart rate variability in depressive and anxiety disorders." *American heart journal* 140, no. 4 (2000): S77-S83.
- [5] Carmigniani, Julie, and Borko Furht. "Augmented reality: an overview." *Handbook of augmented reality* (2011): 3-46.
- [6] Zheng, J. M., K. W. Chan, and Ian Gibson. "Virtual reality." *Ieee Potentials* 17, no. 2 (1998): 20-23.
- [7] Harris, Sandra R., Robert L. Kemmerling, and Max M. North. "Brief virtual reality therapy for public speaking anxiety." *Cyberpsychology & behaviour* 5, no. 6 (2002): 543-550.
- [8] Bekelis, Kimon, Daniel Calnan, Nathan Simmons, Todd A. MacKenzie, and George Kakoulides. "Effect of an immersive preoperative virtual reality experience on patient reported outcomes: a randomised controlled trial." (2017).
- [9] Wang, Rulin, Xin Huang, Yuan Wang, and Masod Akbari. "Non-pharmacologic Approaches in Preoperative Anxiety, a Comprehensive Review." *Frontiers in Public Health* 10 (2022). <https://doi.org/10.3389/fpubh.2022.854673>
- [10] Everyone has heroic potential. SuperBetter. (n.d.). Retrieved January 3, 2023, from <https://www.superbetter.com/>.
- [11] Tgc. (2019, February 14). Flower. thatgamecompany.com/flower/.
- [12] Home. Personal Zen. (2022, September 20). Retrieved January 3, 2023, from <https://personalzen.com/>
- [13] Inal, Sevil, and Meral Kelleci. "Distracting children during blood draw: Looking through distraction cards is effective in pain relief of children during blood draw." *International journal of nursing practice* 18, no. 2 (2012): 210-219. <https://doi.org/10.1111/j.1440-172X.2012.02016.x>
- [14] Vogt, Lina, Martin Klasen, Rolf Rossaint, Ute Goeretz, Peter Ebus, and Sasa Sopka. "Virtual reality tour to reduce preoperative anxiety in an operating setting before anesthesia: Randomized clinical trial." *Journal of medical Internet research* 23, no. 9 (2021): e28018. <https://doi.org/10.2196/28018>
- [15] Mallari B, Spaeth EK, Goh H, Boyd BS. Virtual reality as an analgesic for acute and chronic pain in adults: a systematic review and meta-analysis. *J Pain Res.* 2019 Jul 3;12:2053-2085. <https://doi.org/10.2147/JPR.S200498>

- [16] Julian, Laura J. "Measures of anxiety." *Arthritis care & research* 63, no. 0 11 (2011). <https://doi.org/10.1002/acr.20561>
- [17] Beck, Aaron T., Norman Epstein, Gary Brown, and Robert A. Steer. "An inventory for measuring clinical anxiety: psychometric properties." *Journal of consulting and clinical psychology* 56, no. 6 (1988). <https://doi.org/10.1037/0022-006X.56.6.893>
- [18] Arnold, Lesley M., Daniel Clauw, Fujun Wang, Jonna Ahl, Paula J. Gaynor, and MADELAINE M. Wohlreich. "Flexible dosed duloxetine in the treatment of fibromyalgia: a randomized, double-blind, placebo-controlled trial." *The Journal of rheumatology* 37, no. 12 (2010): 2578-2586. <https://doi.org/10.3899/jrheum.100365>
- [19] Scopaz, Kristen A., Sara R. Piva, Stephen Wisniewski, and G. Kelley Fitzgerald. "Relationships of fear, anxiety, and depression with physical function in patients with knee osteoarthritis." *Archives of physical medicine and rehabilitation* 90, no. 11 (2009): 1866-1873. <https://doi.org/10.1016/j.apmr.2009.06.012>
- [20] Bullock, J. (2021, January 14). Why should I use unity: Top 8 reasons. TurboFuture. Retrieved January 3, 2023, from <https://turbofuture.com/computers/Why-Should-I-Use-Unity-Top-8-Reasons>
- [21] Garbett, S. L. (2022, June 4). Unreal vs. unity for VR game development: Which is best? MUO. Retrieved January 3, 2023, from <https://www.makeuseof.com/unreal-vs-unity-vr-best-game-development/>
- [22] Technologies, U. (n.d.). VR Overview. Unity. Retrieved January 3, 2023, from <https://docs.unity3d.com/540/Documentation/Manual/VROverview.html%20Unity%20VR%20support,this%20for%20each%20build%20target>.
- [23] Chen, Kevin W., Christine C. Berger, Eric Manheimer, Darlene Forde, Jessica Magidson, Laya Dachman, and C. W. Lejuez. "Meditative therapies for reducing anxiety: A systematic review and meta-analysis of randomized controlled trials." *Depression and anxiety* 29, no. 7 (2012): 545-562. <https://doi.org/10.1002/da.21964>
- [24] Cuncic, A. (2020, December 10). The benefits of meditation for generalized anxiety disorder. Verywell Mind. Retrieved January 5, 2023, from <https://www.verywellmind.com/the-benefits-of-meditation-for-generalized-anxiety-disorder-4143127>
- [25] Vive United Kingdom: Discover virtual reality beyond imagination. VIVE. (n.d.). Retrieved February 9, 2023, from <https://www.vive.com/uk/>
- [26] Mayo Foundation for Medical Education and Research. (2018, May 4). Anxiety disorders. Mayo Clinic. Retrieved January 6, 2023, from <https://www.mayoclinic.org/diseases-conditions/anxiety/symptoms-causes/syc-20350961>
- [27] Mekni, Mehdi, and Andre Lemieux. "Augmented reality: Applications, challenges and future trends." *Applied computational science* 20 (2014): 205-214.
- [28] Cabral, Luana L., Fábio Y. Nakamura, Joice MF Stefanello, Luiz CV Pessoa, Bruno PC Smirmaul, and Gleber Pereira. "Initial validity and reliability of the Portuguese Borg rating of perceived exertion 6-20 scale." *Measurement in Physical Education and Exercise Science* 24, no. 2 (2020): 103-114. <https://doi.org/10.1080/1091367X.2019.1710709>
- [29] Spielberger, C. (1983). *Manual for the State-trait Anxiety Inventory (Form Y)*. 121. <https://doi.org/10.1037/t06496-000>
- [30] Farach, Frank J., Larry D. Pruitt, Janie J. Jun, Alissa B. Jerud, Lori A. Zoellner, and Peter P. Roy-Byrne. "Pharmacological treatment of anxiety disorders: Current treatments and future directions." *Journal of anxiety disorders* 26, no. 8 (2012): 833-843.
- [31] Chalmers, John A., Daniel S. Quintana, Maree J-Anne Abbott, and Andrew H. Kemp. "Anxiety disorders are associated with reduced heart rate variability: a meta-analysis." *Frontiers in psychiatry* 5 (2014). <https://doi.org/10.3389/fpsyt.2014.00080>
- [32] Cheng, Ying-Chih, Min-I. Su, Cheng-Wei Liu, Yu-Chen Huang, and Wei-Lieh Huang. "Heart rate variability in patients with anxiety disorders: A systematic review and meta-analysis." *Psychiatry and Clinical Neurosciences* 76, no. 7 (2022): 292-302. <https://doi.org/10.1111/pcn.13356>
- [33] Kazmi, Syed Zaki Hassan, Henggui Zhang, Wajid Aziz, Oliver Monfredi, Syed Ali Abbas, Saeed Arif Shah, Syeda Sobia Hassan Kazmi, and Wasi Haider Butt. "Inverse correlation between heart rate variability and heart rate demonstrated by linear and nonlinear analysis." *PloS one* 11, no. 6 (2016). <https://doi.org/10.1371/journal.pone.0157557>
- [34] Tadesse, Muhiddin, Siraj Ahmed, Teshome Regassa, Timsel Girma, and Ayub Mohammed. "The hemodynamic impacts of preoperative anxiety among patients undergoing elective surgery: An institution-based prospective cohort study." *International Journal of Surgery Open* 43 (2022). <https://doi.org/10.1016/j.ijso.2022.100490>
- [35] Kowal, Magdalena, Eoin Conroy, Niall Ramsbottom, Tim Smithies, Adam Toth, and Mark Campbell. "Gaming your mental health: A narrative review on mitigating symptoms of depression and anxiety using commercial video games." *JMIR serious games* 9, no. 2 (2021): e26575. <https://doi.org/10.2196/26575>
- [36] Stein, Dan J., Naomi A. Fineberg, O. Joseph Bienvenu, Damiaan Denys, Christine Lochner, Gerald Nestadt, James F. Leckman, Scott L. Rauch, and Katharine A. Phillips. "Should OCD be classified as an anxiety disorder in DSM-V?." *Depression and anxiety* 27, no. 6 (2010): 495-506.
- [37] Stein, Murray B., and Jitender Sareen. "Generalized anxiety disorder." *New England Journal of Medicine* 373, no. 21 (2015): 2059-2068.
- [38] McNally, Richard J. "Anxiety sensitivity and panic disorder." *Biological psychiatry* 52, no. 10 (2002): 938-946.
- [39] Norr, Aaron M., Brian J. Albanese, Joseph W. Boffa, Nicole A. Short, and Norman B. Schmidt. "The relationship between gender and PTSD symptoms: Anxiety sensitivity as a mechanism." *Personality and Individual Differences* 90 (2016): 210-213.
- [40] Stein, Murray B., and Dan J. Stein. "Social anxiety disorder." *The lancet* 371, no. 9618 (2008): 1115-1125.